



U.S. Department  
of Transportation  
**Federal Aviation  
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15  
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM  
FAA NTSB COUNSEL

Mike Monroney  
Aeronautical Center

P.O. Box 25082  
Oklahoma City, Oklahoma 73125

Tuesday, January 02, 2018

National Transportation Safety Board  
4760 Oakland Street, Suite 500  
Denver, CO 80239

ACCIDENT # 0241    INDIVIDUAL#: 001    NAME [REDACTED]    MODE: AVIATION  
DATE OF ACCIDENT 11/20/2017    DATE RECEIVED 11/29/2017    PUTREFACTION: Yes  
N # 4179M    NTSB # CEN18FA035    CAMI REF # 201700241001  
LOCATION OF ACCIDENT Electra, TX  
SPECIMENS Bile, Blood (Cavity), Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

### FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood (Cavity)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Vitreous

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in ( ) can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marijuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

>> Naproxen detected in Urine

Russell Lewis, Ph.D., F-ABFT  
Supervisor, Forensic Sciences  
Bioaeronautical Sci. Research Lab  
CAMI, FAA