



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Thursday, July 25, 2019

National Transportation Safety Board
4760 Oakland Street, Suite 500
Denver, CO 80239

ACCIDENT # 0106 INDIVIDUAL#: 001 NAME: [REDACTED] MODE: AVIATION
DATE OF ACCIDENT 05/11/2019 DATE RECEIVED 05/30/2019 PUTREFACTION: Yes
N # 713LA NTSB # CEN19FA140 CAMI REF # 201900106001
LOCATION OF ACCIDENT Eagle Lake, TX
SPECIMENS Blood, Brain, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

SUPPLEMENTAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

- NO CARBOXYHEMOGLOBIN detected in Blood

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

- NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

- 47 (mg/dL, mg/hg) Ethanol detected in Blood #1
- 78 (mg/dL, mg/hg) Ethanol detected in Urine
- 94 (mg/dL, mg/hg) Ethanol detected in Vitreous
- 53 (mg/dL, mg/hg) Ethanol detected in Blood #2
- NO ETHANOL detected in Liver
- 98 (mg/dL, mg/hg) Ethanol detected in Heart
- Propanol (N-) detected in Heart
- 113 (mg/dL, mg/hg) Ethanol detected in Kidney
- Propanol (N-) detected in Kidney
- 54 (mg/dL, mg/hg) Ethanol detected in Spleen

DRUGS: Specimens are analyzed using immunoassay, chromatography, mass spectrometry, or spectrophotometry. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

- NO DRUGS listed above detected in Urine

[REDACTED]

Thursday, July 25, 2019

CONTINUATION OF REF#: 201900106001 — [REDACTED]


[REDACTED]

c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J LEWIS
2019.07.29 15:42:01 -05'00'

Russell Lewis, Ph.D., F-ABFT
Supervisor, Forensic Sciences
Bioaeronautical Sci. Research Lab
CAMI, FAA