

10 oz FOAM BUSTER
62.5 gal BUCCANEER XTRA
300 gal KQ-XRN
87.5 WATER

450 TOTAL SOLUTION



KQ-XRN

28-0-0 with 72% Slow Release Nitrogen TECH SHEET

GUARANTEED ANALYSIS:

Total Nitrogen (N)*	28%
7.8% Urea Nitrogen	
20.2% Slowly available water soluble nitrogen* derived from urea, polymethylene urea, methylene diurea and monomethylol urea.	
20.2% Slowly available water soluble nitrogen* from urea, polymethylene urea, methylene diurea and monomethylol urea.	
Weight per Gallon	10.5 lbs.
Gallons per Ton	187.7
Nitrogen per Gallon	2.97 lbs.

MANUFACTURED BY
KUGLER COMPANY • 209 West 3rd; PO Box 1748
McCook, NE 69001
308 -345 -2280
www.kuglercompany.com

CAUTION: KEEP OUT OF REACH OF CHILDREN. Do not get in eyes, on skin or on clothing. Wear goggles, face shield, or safety glasses and rubber gloves when handling. If eye exposure occurs get medical attention immediately. Harmful if swallowed. Avoid breathing vapor. Protect from freezing. Read entire label before using. Wash contaminated clothing before reuse.

WHAT TO DO IN CASE OF CONTACT:
IF IN EYES: Flush eyes with water for 15 minutes, holding eyelids open. Obtain medical attention if irritation develops.
IF ON SKIN: Wash skin thoroughly with soap and water. Get medical attention if irritation develops.
IF SWALLOWED: Drink large volumes of water and induce vomiting by placing finger at back of throat. Call a physician or poison control center.
IF INHALED: Move to fresh air. Obtain medical attention if irritation develops. If contact is made with spray solution containing pesticide(s), follow the "Statement of Practical Treatment/First Aid" the pesticide(s) label.

GENERAL INFORMATION:

Foliar fertilization is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by most crops. 72% of the total Nitrogen in this product is controlled release.

RECOMMENDED FOLIAR RATES:

ALFALFA: 1-2 gals/acre. Apply after each cutting when

CABBAGE, CAULIFLOWER, BROCCOLI: 2-3 gals/acre at early head formation and repeat 7-10 days later.
CELERY: 1-2 gals/acre 10 days after transplanting. Repeat at 2 gals/acre in 3 intervals until 3 weeks prior to harvest, using 7-8 gallons total over crop.
CITRUS: 1-5 gals/acre at pre-bloom or bloom to increase fruit set and repeat in 30 days. Subsequent applications when nitrogen needs are evident
CORN: May apply up to 5 gals/acre with seed at planting. Foliar apply up to 2 gals/acre any time extra nitrogen is needed.
CORN (SEED): May apply up to 5 gals/acre with seed at planting. Foliar apply 2 gals/acre before detasseling and repeat after pollination.
COTTON: Early Season: 1-2 quarts/acre in an over the row application after plants reach 4 leaf stage. Mid-Season: 2-5 gals/acre at early bloom and repeat as necessary.
CUCUMBERS, SQUASH, PUMPKINS, WATERMELONS, CANTALOUPE: 1-3 gals/acre at early bloom and repeat approximately 4 weeks after bloom application.
GARLIC: Spray 1-2 gals/acre directed over row from 10-12 inches high. Make 3-4 applications.
GRAPES (TABLE): 1-2 gals/acre when nitrogen needs are evident or an increase in canopy is required to reduce sunburn.
GRAPES (SEED PRODUCTION): 1-1½ gals/acre at pre-bloom. 2-3 gals/acre at seed head elongation.
LETTUCE: 2-3 gal./acre after thinning. Repeat 14 days later and again 14 days prior to harvest.
MILO and OTHER SORGHUMS: May apply up to 3 gals/acre with seed at planting. Foliar apply up to 2 gals/acre at any time extra nutrition is needed.
MINT (PEPPERMINT, SPEARMINT): For row or establish mint: 2-3 gals/acre — end of March, early April
3-5 gals/acre—post flame late April, early May. May be combined with fungicide.
2-3 gals/acre—mid June. Combine with pesticide
3-5 gals/acre—mid July. Combine with pesticide
2-3 gals/acre — Post Harvest. Can be combined with other nutrients. Can be combined with crop protection chemicals when properly buffered. KQ-XRN pH is 8.5-10. Apply in sufficient water, minimum of 20 gals/acre to provide good coverage to foliage.
ONIONS: 2 gals/acre when onions reach 6-8 inches repeat during bulb fill, and again in two weeks prior to harvest.
PEACHES, PLUMS, PRUNES, NECTARINES, APRICOTS: 1-

BUCCANEER® 5 Extra

herbicide

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

ACTIVE INGREDIENT:

Glyphosate*, N-(phosphonomethyl)glycine	53.8%
in the form of its isopropylamine salt	46.2%
OTHER INGREDIENTS:	100.0%
TOTAL	

*Contains 648 grams per litre or 5.4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 480 grams per litre or 4 pounds per U.S. gallon of the acid, glyphosate.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Wear long sleeved shirt, long pants, shoes, and socks.
DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

TENKÖZ

Glyphosate	Group	9	Herbicide
------------	-------	---	-----------

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE

CALL CHEMTREC (800) 424-9300

EPA Reg. No. 55467-15

Tenköz, Inc.
1725 Windward Concourse
Suite 410
Alpharetta, GA 30005

65310 092018 10YR18

FOAMBUSTERTM

MAX

MANUFACTURED FOR
HELENA CHEMICAL COMPANY
COLLIERVILLE, TENNESSEE 38017, U.S.A. • 901-761-0050

2.25