



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Wednesday, October 22, 2014

National Transportation Safety Board
45065 Riverside Parkway
Ashburn, VA 20147

ACCIDENT # 0173 INDIVIDUAL#: 001 NAME: ENSELL, ALBERT E., IV MODE: AVIATION
DATE OF ACCIDENT 08/09/2014 DATE RECEIVED 08/19/2014 PUTREFACTION: Yes
N # 7968Z NTSB # ERA14FA377 CAMI REF # 201400173001
LOCATION OF ACCIDENT Peachtree City, GA
SPECIMENS Bile, Blood, Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Vitreous

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marijuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

>> Ibuprofen detected in Urine
>> Lidocaine detected in Urine
>> Oxycodone detected in Urine
>> Oxycodone NOT detected in Blood
>> Oxymorphone detected in Urine
>> Oxymorphone NOT detected in Blood
>> Salicylate detected in Urine

Digitally signed by RUSSELL J LEWIS
Date: 2014.11.03 14:27:24 -06'00'

Wednesday, October 22, 2014

CONTINUATION OF REF#: 201400173001 — ENSELL, ALBERT E., IV



Digitally signed by RUSSELL J LEWIS

Date: 2014.11.03 14:27:37 -06'00'

Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI



U.S. Department
of Transportation
**Federal Aviation
Administration**

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Wednesday, October 22, 2014

National Transportation Safety Board
45065 Riverside Parkway
Ashburn, VA 20147

ACCIDENT #	0173	INDIVIDUAL#:	001	NAME:	ENSELL, ALBERT E., IV	MODE:	AVIATION
DATE OF ACCIDENT	08/09/2014	DATE RECEIVED	08/19/2014	PUTREFACTION:	Yes		
	N # 7968Z	NTSB #	ERA14FA377	CAMI REF #	201400173001		
LOCATION OF ACCIDENT	Peachtree City, GA						
SPECIMENS	Bile, Blood, Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous						

CLINICAL REPORT

CLINICAL: Vitreous and Urine are tested for the presence of glucose with reagent strips and by enzymatic spectrophotometric analysis. Postmortem vitreous glucose levels above 125 mg/dL are considered abnormal and postmortem urine levels above 100 mg/dL are considered abnormal. Hemoglobin A1C is analyzed using a latex immunoagglutination inhibition methodology. Hemoglobin A1C blood levels above 6% are considered abnormal. Urine specimens are defined as "dilute" if the creatinine concentration is < 20 mg/dL and the specific gravity is < 1.003. Concentrations of serotonin metabolites 5-hydroxytryptophol (5-HTOL) and 5-hydroxyindole-3-acetic acid (5-HIAA) are measured by LC/MS. A 5-HTOL/5-HIAA ratio value < 15 pmol/nmol is not consistent with ethanol ingestion, while a ratio value > 15 pmol/nmol is indicative of ethanol ingestion.

>> Glucose NOT detected in Vitreous
>> 135 (mg/dl) Glucose detected in Urine
>> 7.4 (%) Hemoglobin A1C detected in Blood

Digitally signed by RUSSELL J LEWIS
Date: 2014.11.03 14:27:05 -06'00'

Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI