



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Tuesday, March 08, 2016

National Transportation Safety Board
45065 Riverside Parkway
Ashburn, VA 20147

Mission Pilot

ACCIDENT # 0024 INDIVIDUAL#: 002 NAME: [REDACTED] MODE: AVIATION
DATE OF ACCIDENT 02/01/2016 DATE RECEIVED 02/04/2016 PUTREFACTION: No
N # 784CP NTSB # ERA16FA100 CAMI REF # 201600024002
LOCATION OF ACCIDENT Mobile, AL
SPECIMENS Blood (Heart), Heart, Liver, Lung

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NOT PERFORMED

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Liver
>> NO ETHANOL detected in Heart

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marijuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

>> 0.521 (ug/ml, ug/g) Doxylamine detected in Liver
>> 0.129 (ug/ml, ug/g) Doxylamine detected in Heart
>> Ibuprofen detected in Liver

c=US, o=U.S. Government, ou=AMC, ou=AMC,
cn=RUSSELL J LEWIS
2016.03.11 07:37:00 -06'00'

Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Thursday, March 17, 2016

National Transportation Safety Board
45065 Riverside Parkway
Ashburn, VA 20147

Mission Scanner

ACCIDENT # 0024 INDIVIDUAL#: 001 NAME: [REDACTED] MODE: AVIATION
DATE OF ACCIDENT 02/01/2016 DATE RECEIVED 02/04/2016 PUTREFACTION: No
N # 784CP NTSB # ERA16FA100 CAMI REF # 201600024001
LOCATION OF ACCIDENT Mobile, AL
SPECIMENS Blood, Brain, Heart, Liver, Lung, Muscle, Spleen

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NOT PERFORMED

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Muscle
>> NO ETHANOL detected in Brain

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marijuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

>> Amlodipine detected in Muscle
>> Amlodipine detected in Liver
>> Diphenhydramine detected in Liver
>> Diphenhydramine detected in Muscle
>> Losartan detected in Liver
>> 1.298 (ug/mL, ug/g) Trazodone detected in Liver
>> 0.282 (ug/mL, ug/g) Trazodone detected in Muscle

Thursday, March 17, 2016

CONTINUATION OF REF#: 201600024001 — Dryden, Phil J.



Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI

c=US, o=U.S. Government, ou=AMC, ou=AMC,
cn=RUSSELL J LEWIS
2016.03.23 10:55:31 -05'00'