



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P O Box 25082
Oklahoma City, Oklahoma 73125

Thursday, April 4, 2019

National Transportation Safety Board
45065 Riverside Parkway
Ashburn, VA 20147

ACCIDENT # 0035 INDIVIDUAL#: 001 NAME: [REDACTED] MODE: AVIATION
DATE OF ACCIDENT 02/23/2019 DATE RECEIVED 02/26/2019 PUTREFACTION: No
N # 224TA NTSB # ERA19FA107 CAMI REF # 201900035001
LOCATION OF ACCIDENT Mansfield, MA
SPECIMENS Bile, Blood (Cavity), Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spinal Fluid, Spleen, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

- NO CARBOXYHEMOGLOBIN detected in Blood (Cavity)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

- NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

- NO ETHANOL detected in Blood (Cavity)

DRUGS: Specimens are analyzed using immunoassay, chromatography, mass spectrometry, or spectrophotometry. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

- NO DRUGS listed above detected in Urine

[REDACTED]

c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J LEWIS
2019.04.10 12:44:14 -05'00'

Russell Lewis, Ph.D., F-ABFT
Supervisor, Forensic Sciences
Bioaeronautical Sci. Research Lab
CAMI, FAA



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P O Box 25082
Oklahoma City, Oklahoma 73125

Thursday, April 4, 2019

National Transportation Safety Board
45065 Riverside Parkway
Ashburn, VA 20147

ACCIDENT # 0035 INDIVIDUAL#: 002 NAME: [REDACTED] MODE: AVIATION
DATE OF ACCIDENT 02/23/2019 DATE RECEIVED 02/26/2019 PUTREFACTION: No
N # 224TA NTSB # ERA19FA107 CAMI REF # 201900035002
LOCATION OF ACCIDENT Mansfield, MA
SPECIMENS Bile, Blood, Blood (Cavity), Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

- NO CARBOXYHEMOGLOBIN detected in Blood (Cavity)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.


- NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

- NO ETHANOL detected in Blood (Cavity)

DRUGS: Specimens are analyzed using immunoassay, chromatography, mass spectrometry, or spectrophotometry. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

- NO DRUGS listed above detected in Urine


Russell Lewis, Ph.D., F-ABFT
Supervisor, Forensic Sciences
Bioaeronautical Sci. Research Lab
CAMI, FAA

c=US, o=U.S. Government, ou=AMC, ou=AMC,
cn=RUSSELL J LEWIS
2019.04.10 12:46:42 -05'00'