



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Thursday, September 5, 2019

National Transportation Safety Board
45065 Riverside Parkway
Ashburn, VA 20147

ACCIDENT # 0174 INDIVIDUAL#: 001 NAME: [REDACTED] **Pilot** MODE: AVIATION
DATE OF ACCIDENT 08/11/2019 DATE RECEIVED 08/14/2019 PUTREFACTION: No
N # 43032 NTSB # ERA19FA243 CAMI REF # 201900174001
LOCATION OF ACCIDENT Jersey Shore, PA
SPECIMENS Bile, Blood, Blood (Cavity), Brain, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

- NO CARBOXYHEMOGLOBIN detected in Blood (Cavity)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

- NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

- NO ETHANOL detected in Blood (Cavity)

DRUGS: Specimens are analyzed using immunoassay, chromatography, mass spectrometry, or spectrophotometry. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

- NO DRUGS listed above detected in Urine

[REDACTED]

c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J LEWIS
2019.09.10 15:19:19 -05'00'

Russell Lewis, Ph.D., F-ABFT
Supervisor, Forensic Sciences
Bioaeronautical Sci. Research Lab
CAMI, FAA



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Thursday, September 5, 2019

National Transportation Safety Board
45065 Riverside Parkway
Ashburn, VA 20147

ACCIDENT # 0174 INDIVIDUAL#: 001 NAME: [REDACTED] Pilot MODE: AVIATION
DATE OF ACCIDENT 08/11/2019 DATE RECEIVED 08/14/2019 PUTREFACTION: No
N # 43032 NTSB # ERA19FA243 CAMI REF # 201900174001
LOCATION OF ACCIDENT Jersey Shore, PA
SPECIMENS Bile, Blood, Blood (Cavity), Brain, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

CLINICAL REPORT

CLINICAL: Vitreous and Urine are tested for the presence of glucose with reagent strips and by enzymatic spectrophotometric analysis. Postmortem vitreous glucose levels above 125 mg/dL are considered abnormal and postmortem urine levels above 100 mg/dL are considered abnormal. Hemoglobin A1C is analyzed using a latex immunoagglutination inhibition methodology. Hemoglobin A1C blood levels above 6% are considered abnormal. Urine specimens are defined as "dilute" if the creatinine concentration is < 20 mg/dL and the specific gravity is < 1.003. Concentrations of serotonin metabolites 5-hydroxytryptophol (5-HTOL) and 5-hydroxyindole-3-acetic acid (5-HIAA) are measured by LC/MS. A 5-HTOL/5-HIAA ratio value < 15 pmol/nmol is not consistent with ethanol ingestion, while a ratio value > 15 pmol/nmol is indicative of ethanol ingestion.

- 26 (mg/dL) Glucose detected in Vitreous
- 1910 (mg/dL) Glucose detected in Urine
- 7.8 (%) Hemoglobin A1C detected in Blood (Cavity)

[REDACTED]
Russell Lewis, Ph.D., F-ABFT
Supervisor, Forensic Sciences
Bioaeronautical Sci. Research Lab
CAMI, FAA

c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J
LEWIS
2019.09.10 15:19:42 -05'00'