



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Wednesday, October 18, 2017

National Transportation Safety Board
505 South 336th Street, Suite 540
Federal Way, WA 98003

ACCIDENT #	0203	INDIVIDUAL#:	001	NAME:	[REDACTED]	MODE:	AVIATION
DATE OF ACCIDENT	09/22/2017	DATE RECEIVED	09/29/2017	PUTREFACTION:	No		
	N # 7070Q	NTSB #	WPR17FA213	CAMI REF #	201700203001		
LOCATION OF ACCIDENT	Perkinsville, AZ						
SPECIMENS	Bile, Blood (Heart), Brain, Gastric, Heart, Kidney, Liver, Lung, Spleen						

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood (Heart)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Blood (Heart)

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marijuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

>> NO DRUGS listed above detected in Blood (Heart)

[REDACTED]
Russell Lewis, Ph.D., F-ABFT
Supervisor, Forensic Sciences
Bioaeronautical Sci. Research Lab
CAMI, FAA

c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J LEWIS
2017.10.23 14:30:26 -05'00'



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Wednesday, October 18, 2017

National Transportation Safety Board
505 South 336th Street, Suite 540
Federal Way, WA 98003

ACCIDENT # 0203 INDIVIDUAL#: 002 NAME: [REDACTED] MODE: AVIATION
DATE OF ACCIDENT 09/22/2017 DATE RECEIVED 09/29/2017 PUTREFACTION: No
N # 7070Q NTSB # WPR17FA213 CAMI REF # 201700203002
LOCATION OF ACCIDENT Perkinsville, AZ
SPECIMENS Bile, Blood (Cardiac), Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood (Cardiac)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Vitreous

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marijuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

>> NO DRUGS listed above detected in Urine

[REDACTED]

c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J LEWIS
2017.10.23 14:31:10 -05'00'

Russell Lewis, Ph.D., F-ABFT
Supervisor, Forensic Sciences
Bioaeronautical Sci. Research Lab
CAMI, FAA