

BASE LABEL TEXT

DANGER**KEEP OUT OF REACH OF CHILDREN**

The Syngenta logo, which consists of the word "syngenta" in a white, lowercase, sans-serif font. A small registered trademark symbol (®) is located at the end of the word. The logo is set against a green rectangular background.

Active Ingredient:

BRAVO® WEATHER STIK® contains 720 g/litre CHLOROTHALONIL in the form of a suspension concentrate.

GROUP	M5	FUNGICIDE
-------	----	-----------

For the control of fungal diseases in Fruit, Ornamentals, Turf, Vegetables and Wheat.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

Net Contents: 10 litres

*Registered pursuant to the ACVM Act 1997, No. P7065.
See www.foodsafety.govt.nz for registration conditions*

Approved pursuant to the HSNO Act 1996, Approval Code HSR000670.

Syngenta Crop Protection Limited
Level 4, 60 Parnell Road, Parnell, Auckland 1052

®

PLEASE READ THE LABEL COMPLETELY BEFORE USE

This product **MUST NOT** be used for any purpose or in any manner contrary to the controls and conditions of this registration.

This product **MUST** be Tracked.

Any person purchasing, handling or disposing of this product must hold a Certified Handler certificate.

Any person mixing, loading or handling this product must be a qualified loader.

Any person applying this product must be a qualified contractor or under the guidance of a qualified contractor.

This substance is restricted to use in the workplace only.

**GHS Classifications**

Acute inhalation toxicity Category 2

Serious eye damage Category 1

Skin sensitisation Category 1

Carcinogenicity Category 2

Specific target organ toxicity (repeated exposure) Category 1

Hazardous to the aquatic environment acute Category 1

Hazardous to the aquatic environment chronic Category 1

Hazardous to soil organisms

Hazardous to terrestrial vertebrates

DANGER

Fatal if inhaled.

Causes serious eye damage

May cause an allergic skin reaction.

Suspected of causing cancer.

Causes kidney damage from repeated oral exposure at high doses.

Very toxic to aquatic life with long lasting effects.

Harmful to the soil environment.

Toxic to terrestrial vertebrates.

PRECAUTIONS

Store in original container, tightly closed, away from foodstuffs and under lock and key.

When handling, wear full protective clothing such as overalls, chemical resistant gloves, respirator and face shield.

Wash splashes of concentrate from skin or eyes immediately.

When spraying, wear overalls and chemical resistant gloves.

Avoid contact with skin, eyes and inhalation of spray mist.

DO NOT eat, drink or smoke while using.

Wash hands and face before eating, drinking or smoking and after work.

Wash protective clothing after use.

Apply with well maintained and calibrated spray equipment.

Tracking records **MUST** be kept as per schedule 27 of Health and Safety at Work (Hazardous Substances) Regulations 2017.

Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.

DO NOT carry on a passenger service vehicle.

Avoid contamination of any water supply with chemical or empty container.

Container disposal: Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.

Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility. For additional information refer to EPA Notices: Hazardous Substance (Disposal) Notice 2017.

FIRST AID

If SWALLOWED, DO NOT induce vomiting. Rinse Mouth.

If SKIN or HAIR contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin. If skin irritation or rash occurs get medical advice.

If splashed in the EYES, wash out immediately with water. Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor. If eye irritation persists: Get medical advice/attention.

If INHALED, move the victim to fresh air and keep at rest in a position comfortable for breathing.

Immediately call a POISON CENTER or doctor. Specific treatment is urgent.

Begin artificial respiration if breathing has stopped. Use mouth to nose rather than mouth to mouth.

Obtain medical attention. Take to hospital without delay.

For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If medical advice is needed, have product container or label at hand.

SPILLAGE

In the event of a leak or spillage - contain the spillage, taking particular care to prevent entry to drains or waterways:

- Keep people away.
- Absorb spillage with material such as sawdust or soil, and dispose of safely.
- Sweep up contaminated material and place in drums for disposal in accordance with the requirements of the EPA New Zealand controls and in a manner acceptable to the local authorities. For major spillages, inform the Fire Service immediately.
- Overalls, eye protection, chemical resistant gloves and gumboots should be worn during clean up and disposal.

STORAGE

Store in the closed, original container in a cool, dry, well ventilated area. DO NOT store for prolonged periods in direct sunlight. See the Safety Data Sheet for additional storage information. Stores containing quantities of 100 litres or more require signage and secondary containment.

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



Distributed by and registered to Syngenta Crop Protection Limited, Level 4, 60 Parnell Road, Parnell, Auckland. Phone 0800 333 336.

Batch No	
Date of manufacture	

BARCODE

UN 3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.
(chlorothalonil) MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2X
CONTAIN SPILLAGE



In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 734 607 (24 hours).

LABEL BOOKLET TEXT

DANGER**KEEP OUT OF REACH OF CHILDREN**

The Syngenta logo, which consists of the word "syngenta" in a white, lowercase, sans-serif font. A small registered trademark symbol (®) is located to the upper right of the "a". The logo is set against a green rectangular background.

Active Ingredient:

BRAVO® WEATHER STIK® contains 720 g/litre CHLOROTHALONIL in the form of a suspension concentrate.

GROUP	M5	FUNGICIDE
--------------	-----------	------------------

For the control of fungal diseases in Fruit, Ornamentals, Turf, Vegetables and Wheat.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

Net Contents: 10 litres

*Registered pursuant to the ACVM Act 1997, No. P7065.
See www.foodsafety.govt.nz for registration conditions*

Approved pursuant to the HSNO Act 1996, Approval Code HSR000670.

Syngenta Crop Protection Limited

Level 4, 60 Parnell Road, Parnell, Auckland 1052

®

PLEASE READ THE LABEL COMPLETELY BEFORE USE

This product **MUST NOT** be used for any purpose or in any manner contrary to the controls and conditions of this registration.

This product **MUST** be Tracked.

Any person purchasing, handling or disposing of this product must hold a Certified Handler certificate.

Any person mixing, loading or handling this product must be a qualified loader.

Any person applying this product must be a qualified contractor or under the guidance of a qualified contractor.

This substance is restricted to use in the workplace only.

**GHS Classifications**

Acute inhalation toxicity Category 2

Serious eye damage Category 1

Skin sensitisation Category 1

Carcinogenicity Category 2

Specific target organ toxicity (repeated exposure) Category 1

Hazardous to the aquatic environment acute Category 1

Hazardous to the aquatic environment chronic Category 1

Hazardous to soil organisms

Hazardous to terrestrial vertebrates

DANGER

Fatal if inhaled.

Causes serious eye damage

May cause an allergic skin reaction.

Suspected of causing cancer.

Causes kidney damage from repeated oral exposure at high doses.

Very toxic to aquatic life with long lasting effects.

Harmful to the soil environment.

Toxic to terrestrial vertebrates..

PRECAUTIONS

Store in original container, tightly closed, away from foodstuffs and under lock and key.

When handling, wear full protective clothing such as overalls, chemical resistant gloves, respirator and face shield.

Wash splashes of concentrate from skin or eyes immediately.

When spraying, wear overalls and chemical resistant gloves.

Avoid contact with skin, eyes and inhalation of spray mist.

DO NOT eat, drink or smoke while using.

Wash hands and face before eating, drinking or smoking and after work.

Wash protective clothing after use.

Apply with well maintained and calibrated spray equipment.

Tracking records **MUST** be kept as per schedule 27 of Health and Safety at Work (Hazardous Substances) Regulations 2017.

Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.

DO NOT carry on a passenger service vehicle.

Avoid contamination of any water supply with chemical or empty container.

Container disposal: Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.

Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility. For additional information refer to EPA Notices: Hazardous Substance (Disposal) Notice 2017.

FIRST AID

If SWALLOWED, DO NOT induce vomiting. Rinse Mouth.

If SKIN or HAIR contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin. If skin irritation or rash occurs get medical advice.

If splashed in the EYES, wash out immediately with water. Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor. If eye irritation persists get medical advice/attention.

If INHALED, move the victim to fresh air and keep at rest in a position comfortable for breathing.

Immediately call a POISON CENTER or doctor. Specific treatment is urgent.

Begin artificial respiration if breathing has stopped. Use mouth to nose rather than mouth to mouth.

Obtain medical attention. Take to hospital without delay.

For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If medical advice is needed, have product container or label at hand.

SPILLAGE

In the event of a leak or spillage - contain the spillage, taking particular care to prevent entry to drains or waterways:

- Keep people away.
- Absorb spillage with material such as sawdust or soil, and dispose of safely.
- Sweep up contaminated material and place in drums for disposal in accordance with the requirements of the EPA New Zealand controls and in a manner acceptable to the local authorities. For major spillages, inform the Fire Service immediately.
- Overalls, eye protection, chemical resistant gloves and gumboots should be worn during clean up and disposal.

DIRECTIONS FOR USE

MANDATORY DIRECTIONS

By law, the following restrictions must be complied with:

As BRAVO® WEATHER STIK® contains minute quantities of Hexachlorobenzene (HCB), a compound that can be taken up by grazing stock to give unacceptable residues in meat and milk, the following restrictions on the use of the product **MUST** be observed:

- **DO NOT** graze treated crops;
- **DO NOT** feed any part of a treated crop to stock.

These two restrictions **DO NOT** apply if either of the following is first undertaken:

- **CULTIVATE** treated areas thoroughly (ie **DO NOT** use direct drilling or other minimum tillage techniques) before sowing pasture or any animal feed crop, or
- **SOIL TESTING** in compliance with the '*Guideline for Management of Sheep Grazing in Vineyards or Orchards Contaminated with Hexachlorobenzene*' of the vineyard/orchard area indicates HCB residues are less than 0.001 mg/kg in the soil.

It is an offence to use this product on animals.

Crop	Disease	Rate/100 litres water	Rate/ha	Critical Comments
Asparagus	Stemphyllium Leaf Spot	210 mL	2.1 litres	Commence spraying at first appearance of disease after closing up and repeat at 14 to 28 day intervals to maintain disease control until ferns die off. Use the shorter spray interval when disease pressure is high. Ensure good plant coverage.
Beans Butter, Green, Pole	Rust	175 to 210 mL	1.4 litres	Apply at seedling emergence and continue at 10 to 21 day intervals.
Beans Broad		210 to 245 mL	2.1 litres	Apply at first appearance of foliar infection and repeat at 10 to 21 day intervals or as needed to maintain disease control.
Berryfruit Blackberries, Boysenberries, Loganberries	Dryberry, *Botrytis	210 to 245 mL	2.1 litres	Apply as a full season cover spray at 14 day intervals from bud burst onwards. When infection is severe reduce the interval to 7 to 10 days. Autumn applications are necessary to protect primocanes from infection.
Blackcurrants	Leaf Spot	210 mL	2.1 litres	Apply from bud burst onwards at 10 to 14 day intervals.
Brassicas Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Downy Mildew, Ring Spot	175 to 210 mL	1.4 litres	Apply at seedling emergence and continue at 10 to 21 day intervals. Use the high rate for Ring Spot control.
Celery	Septoria Leaf Spot	210 to 245 mL	2.1 litres	Apply as a full season cover spray from immediately post-planting at 7 to 14 day intervals.
Cucurbits Cucumbers, Melons, Squash	Downy Mildew, Powdery Mildew	175 to 210 mL	1.4 litres	Apply at seedling emergence and continue at 10 to 21 day intervals.
Lettuce	Ring Spot	210 to 245 mL	2.1 litres	Apply at seedling emergence and continue at 7 to 10 day intervals throughout the growing period. A regular protective program is necessary to prevent disease establishment.
Ornamentals Camellias, Carnations, Chrysanthemums, Geraniums, Statice, Roses	Black Spot, *Botrytis, Downy Mildew, Flower Blight, Powdery Mildew	175 to 210 mL	-	Apply from the commencement of new growth at 10 to 21 day intervals. DO NOT spray when flower buds are showing petal colour as bloom damage may occur.

Crop	Disease	Rate/100 litres water	Rate/ha	Critical Comments
Pipfruit non bearing Apples and Pears	Black Spot	210 to 245 mL	-	Apply as a full season cover spray at 14 to 21 day intervals. DO NOT apply to bearing trees as fruit russet may occur.
Potatoes	Early Blight, Late Blight	140 mL	1.1 litres	Apply at 7 to 14 day intervals or as needed to maintain disease control.
Stonefruit Apricots, Cherries, Nectarines	Leaf Curl	160 to 210 mL	-	Apply pre bud swell. Where disease levels are high repeat application from pink to shuck fall at 7 to 14 day intervals.
Stonefruit Peaches, Plums	Brown Rot, Rust, Blossom Blight			Apply from pink to shuck fall at 7 to 14 day intervals.
Strawberries	Alternaria Leaf Spot, *Botrytis	210 to 245 mL	2.1 litres	Apply as a full season cover spray at 14 day intervals from bud burst onwards. When infection is severe reduce the interval to 7 to 10 days.
Tomatoes Outdoor	Early Blight, Late Blight, *Botrytis			Apply at planting out and continue at 10 to 14 day intervals until harvest.
Turf, Cotula	Dollar Spot, Brown Patch, Leaf Spotting/ Melting Out	70 to 140 mL/ 100 square metres	-	Spray in sufficient water to thoroughly wet foliage (7.5 to 10 litres per 100 square metres). Use the low rate for preventative applications and the high rate as a curative application. Under conditions of high pressure repeat at 10 to 14 day intervals. As the surface of diseased turf may be hard to wet, especially where thatch is present, add a wetting agent to ensure adequate penetration and control of residual infection resources. MUST NOT be used on areas to be grazed.
	Anthraxnose, Fusarium Patch, Red Thread	105 to 210 mL/ 100 square metres		
Wheat DO NOT use if stock are likely to graze the stubble DO NOT remove the straw from treated fields	<i>Ascochyta</i> spp, Didymella, Glume Blotch, Sooty Moulds	-	0.7 to 1.4 litres	Apply at flag leaf emergence and again up until ear fully emerged if cool, wet conditions prevail. Use the higher rate where disease risk levels are high. Tank mix BRAVO® WEATHER STIK® with other cereal fungicides for control of other diseases.
*Botrytis: BRAVO® WEATHER STIK® applied as a full season cover spray will prevent Botrytis. However, during periods of high risk (wet, humid weather) use a specific Botrytis fungicide.				

GENERAL INFORMATION

BRAVO® WEATHER STIK® is a protectant fungicide that will provide effective control of fungus diseases on a wide range of vegetable, fruit, ornamental crops, turf and wheat. BRAVO® WEATHER STIK® can be used effectively through high volume, low volume or aircraft spray equipment.

Note: BRAVO® WEATHER STIK® must be used only as directed on this label.

RESISTANCE MANAGEMENT

GROUP M5 FUNGICIDE

BRAVO® WEATHER STIK® Fungicide contains A member of the CHLORONITRILE group of fungicides. BRAVO® WEATHER STIK® is a Group M5 fungicide. Resistance to this fungicide could develop from repeated use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies. Since the occurrence of resistant fungi is difficult to detect prior to use, Syngenta Crop Protection Ltd accepts no liability for any losses that may result from the failure of BRAVO® WEATHER STIK® to control resistant fungi. Consult your local supplier, consultant, a Syngenta representative or New Zealand Plant Protection Society (Inc), www.resistance.nzpps.org for alternate modes of action and the details of resistance management strategies for the crops listed on the label.

MIXING

If the container has been standing for some while, roll the drum from side to side several times to ensure a uniform mixture is drawn off. Add the required amount of BRAVO® WEATHER STIK® to the partly filled spray tank and top up with water. Keep mechanical agitator or bypass running while filling, during spraying, and before recommencing spraying after a stoppage.

APPLICATION

Suggested spray volume: It is important that good spray coverage of the crop is achieved and for high volume spraying use a minimum of 800 litres of water per hectare, increasing as necessary to ensure complete foliar cover at all stages of crop development. The rates suggested are for high volume spraying to run off. For concentrate spraying adjust dilution rates accordingly.

SPRAY DRIFT

Take all reasonable steps to ensure that BRAVO® WEATHER STIK® does NOT cause any adverse effects to the environment beyond the application area.

COMPATIBILITY

BRAVO® WEATHER STIK® is compatible with most commonly used fungicides and insecticides. Unless prior experience has shown the combination to be physically compatible and non-injurious to crop, DO NOT combine with emulsifiable formulations, including oils, in the spray tank. DO NOT use with lime sulphur or Bordeaux mixture. BRAVO® WEATHER STIK® is compatible with streptomycin: when using add BRAVO® WEATHER STIK® to the spray tank, top up, then add streptomycin containing product.

WETTING AGENTS

BRAVO® WEATHER STIK® is formulated with an effective wetting system, and the addition of further surfactant is neither required, nor recommended except where suggested in the Directions for Use table.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

Beans (Broad, Butter, Green, Pole)	7 days
Berryfruit (Boysenberries, Blackberries, Loganberries)	7 days
Blackcurrants	7 days
Brassicas (Broccoli, Brussels Sprouts, Cabbage, Cauliflower)	7 days
Celery	7 days
Cucurbits (Cucumbers, Melons, Squash)	1 day
Lettuce	10 days
Potatoes	7 days
Tomatoes (Outdoors)	7 days
Strawberries	1 day
Stonefruit (Apricots, Cherries, Nectarines, Peaches, Plums)	DO NOT use after shuckfall
Wheat	56 days

STORAGE

Store in the closed, original container in a cool, dry, well ventilated area. DO NOT store for prolonged periods in direct sunlight. See Safety Data Sheet for additional storage information. Stores containing quantities of 100 litres or more require signage and secondary containment.

SAFETY DATA SHEET

If additional information is required, refer to the Safety Data Sheet. For a copy visit our website at www.syngenta.co.nz

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



Distributed by and registered to Syngenta Crop Protection Limited, Level 4, 60 Parnell Road, Parnell, Auckland 1052. Phone 0800 333 336.



WILBUR-ELLIS®

Ideas to Grow With

GUARANTEED ANALYSIS

Total Nitrogen (N)	9.00%
1.00% Nitrate Nitrogen	
3.30% Urea Nitrogen	
4.70% Other Water Soluble Nitrogen*	
Calcium (Ca)	1.35%
DERIVED FROM: Low Biuret Urea, Triazone, Methylene Urea, Calcium Nitrate, and Barley Protein Hydrolysate.	
Biuret (Max.)	0.02%

*4.60% Slowly available nitrogen from triazone and methylene urea.

KEEP OUT OF REACH OF CHILDREN PRECAUTION

Do not ingest. Avoid contact with skin, eyes or clothing.
Avoid breathing dust, vapor or mist.

FIRST AID

In all cases, call a poison control center or doctor for further treatment advice.

IF SWALLOWED, call a poison control center or doctor immediately. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. **IF ON SKIN**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF INHALED**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration. **IF IN EYES**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.

For chemical spills, leaks, fire or exposure,
call CHEMTREC: (800) 424-9300.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Store in a safe manner. Store in original container only and keep tightly sealed when not in use. Dispose of unused product and empty containers in accordance with federal, state and local regulations.

GENERAL INFORMATION

FOLI-GRO® GR BOOSTER PLUS is formulated primarily for foliar applications to prevent or correct nitrogen deficiencies in a wide range of agronomic and ornamental plants. Its use is suggested as a supplement to a regular, balanced fertilizer program to enhance yields and improve quality.

Application of FOLI-GRO GR BOOSTER PLUS is a means of obtaining a quick response to needed elements. Foliar applications of FOLI-GRO GR BOOSTER PLUS may be particularly beneficial during periods of peak nutrient demand, for crops grown on soils having poor nutrient availability or to crops suffering from a weakened root system.

Avoid foliar nutrient applications to crops during periods of moisture, salinity or heat stress. Applications of foliar nutrients to sensitive crops such as leafy vegetables or soft-skinned fruit during periods of excessive temperatures may cause cosmetic injury. FOLI-GRO GR BOOSTER PLUS promotes stronger, more vigorous budding and healthier trees. Also helps improve fertilizer efficiency and increases nut size.

FOLI-GRO GR BOOSTER PLUS should be used as part of a comprehensive Total Nutrition System® for optimizing plant growth, development, yield and quality.

DIRECTIONS FOR USE

FOLI-GRO GR BOOSTER PLUS may be applied as a foliar spray by air or by ground. Maintain agitation to provide uniform distribution within the spray tank. Thorough and even spray coverage is essential for maximum nutrient uptake and crop safety. Always include a high quality wetting agent, such as R-11® or SYL-COAT®, at recommended rates.

Optimum rate of application will vary based on a variety of factors including degree of need or deficiency, growth stage, yield potential, environmental conditions, existing soil fertility levels, etc.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aaplco.org/metals.html>

TREE FRUIT AND NUT CROPS: Spring: Make 3 applications at 1 gallon per acre beginning at pink tip and continuing through May. **Fall and Spring:** Apply 1 gallon per acre in the fall, post-harvest but before leaves fall; follow with 2 applications in the spring at 1 gallon per acre, starting at pink tip and continuing through May.

CUCURBITS, BEANS AND FRUITING VEGETABLES: Apply 2-4 quarts per acre during the period of early bloom through early fruit set. An additional application of 2-4 quarts per acre during fruit sizing may be needed.

GRAPES: Apply 1 gallon of FOLI-GRO GR BOOSTER PLUS per acre during the period of bloom to fruit set and again during the period of veraison to maturation.

COMPATIBILITY

FOLI-GRO GR BOOSTER PLUS is compatible with many co-applied nutrient sources as well as plant protection chemicals. FOLI-GRO GR BOOSTER PLUS is compatible with zinc sulfate applications and bloom fungicide sprays. Always jar test new combinations for compatibility prior to field mixing.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of the product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Buyer and User accept all risks arising from any use of this product, including without limitation uses contrary to label instructions, under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

To the extent permitted by law, neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS logo, Ideas to Grow With, FOLI-GRO, R-11, SYL-COAT, and Total Nutrition System are registered trademarks, and FOLI-GRO logo is a trademark of Wilbur-Ellis Company LLC.

K-052318

Manufactured by: WILBUR-ELLIS COMPANY LLC
7 E. Washington Ave., Yakima, WA 98903

LOT#

NET CONTENTS: 2.5 GALLONS (9.46 Liters)

WEIGHT PER GALLON: 9.5 POUNDS at 68°F